



CARDI·OH

Ohio Cardiovascular and Diabetes Health Collaborative



In partnership with:



Cardi-OH ECHO

Health Equity and Cardiovascular Risk

April 11, 2024

About Cardi-OH

Founded in 2017, the mission of Cardi-OH is to improve cardiovascular and diabetes health outcomes and eliminate disparities in Ohio's Medicaid population.

WHO WE ARE: An initiative of health care professionals across Ohio's seven medical schools.

WHAT WE DO: Identify, produce, and disseminate evidence-based cardiovascular and diabetes best practices to primary care teams.

HOW WE DO IT: Best practices resources are available via an online library at Cardi-OH.org, including monthly newsletters, podcasts, webinars, and virtual clinics using the Project ECHO® virtual training model.

[Learn more at Cardi-OH.org](http://Cardi-OH.org)



CARDI•OH

Ohio Cardiovascular and Diabetes Health Collaborative



Cardi-OH ECHO Team

FACILITATOR

Goutham Rao, MD, FAHA
Case Western Reserve University

CONTENT EXPERTS

Karen Bailey, MS, RDN, LD, CDCES
Ohio University

Kristen Berg, PhD
Case Western Reserve University

Elizabeth Beverly, PhD
Ohio University

Danette Conklin, PhD
Case Western Reserve University

Kathleen Dungan, MD, MPH
The Ohio State University

Adam Perzynski, PhD
Case Western Reserve University

Marilee Clemons, PharmD
University of Toledo

Chris Taylor, PhD
The Ohio State University

Kelsey Ufholz, PhD
Case Western Reserve University

James Werner, PhD, MSSA
Case Western Reserve University

Jackson Wright, MD, PhD
Case Western Reserve University

Disclosure Statements



- The following speakers and subject matter experts have a relevant financial interest or affiliation with one or more organizations that could be perceived as a real or apparent conflict of interest in the context of the subject of their presentation*:
 - Danette Conklin, PhD; Kathleen Dungan, MD, MPH; Adam T. Perzynski, PhD; Christopher A. Taylor, PhD, RDN, LD, FAND; Jackson Wright, MD, PhD
- The remaining speakers and subject matter experts have no financial relationships with any commercial interest related to the content of this activity:
 - Karen Bailey, MS, RDN, LD, CDCES; Kristen Berg, PhD; Elizabeth Beverly, PhD; Merilee Clemons, PharmD; Revital Gordodeski Baskin, MD; Allyson Hughes, PhD; George Matar, MD; Kelsey Ufholz, PhD; Goutham Rao, MD, FAHA; James Werner, PhD, MSSA
- The following members of the planning committee DO NOT have any disclosures/financial relationships from any ineligible companies:
 - Shari Bolen, MD; Anderson Christopher; Richard Cornachione; Carolyn Henceroth; Gillian Irwin; Michael Konstan, MD; Elizabeth Littman; Devin O'Neill; Steven Ostrolencki; Ann Nevar; Claire Rollins; Catherine Sullivan

* These financial relationships are outside the presented work.

** For more information about exemptions or details, see www.acme.org/standards



Health Literacy and Cardiovascular Risk

Karen Bailey, MS, RDN, LD, CDCES

Ohio University Diabetes Institute

Learning Objectives



- 1) Describe disparities in health literacy which impact cardiovascular risks, including diabetes
- 2) Describe a culturally sensitive approach to diabetes education
- 3) List strategies to effectively convey numerical information to patients with low health literacy

Health Literacy



Defined as, “the degree to which individuals have the capacity to obtain, process and understand basic health information and services needed to make appropriate health decisions.”

Skills needed include:

- Reading
- Writing
- Calculating numbers
- Communicating with health professionals
- Using health technology (glucose meters, CGMs, connected apps, b/p monitors, etc.)

Estimated that 90 million Americans have low literacy skills.

National Assessment of Adult Literacy (NAAS) - 2003




- Assessed 19,000 adults (16 years and older) in households and prisons.
- Assessed literacy
 - Prose literacy – use information from texts (sentences/paragraphs)
 - Document literacy – use information from noncontinuous texts
 - Quantitative literacy – perform computations using numbers from print text
- Health literacy
 - Health tasks related to clinical information
 - Health tasks related to prevention
 - Health tasks related to navigating the health system

Kutner M, Greenberg E, Jin Y, Paulsen C (2006). The Health Literacy of America's Adults: Results from the 2003 National Assessment of Adult Literacy (NCES 2006-483). U.S. Department of Education. Washington DC: National Center for Education Statistics

Health Literacy Levels

NAAL 2003

Kutner M, Greenberg E, Jin Y, Paulsen C (2006). The Health Literacy of America's Adults: Results from the 2003 National Assessment of Adult Literacy (NCES 2006-483). U.S. Department of Education. Washington DC: National Center for Education Statistics



NAAS – 2003



Health Literacy Levels

- Below Basic
- Basic
- Intermediate
- Proficient

Results

- 53% Americans had intermediate health literacy
- 12% proficient
- 22% basic
- 14% below basic

Health Literacy in US - Results



- Individuals more likely to have low health literacy:
 - Lower socioeconomic status/living below poverty level
 - Lower education level
 - ≥ 65 y/o
 - Non-native speakers of English/low English proficiency
 - Medicare and Medicaid recipients
- Those with low literacy more likely to get health information from radio and TV, not printed materials.

Kutner M, Greenberg E, Jin Y, Paulsen C (2006). The Health Literacy of America's Adults: Results from the 2003 National Assessment of Adult Literacy (NCES 2006-483). U.S. Department of Education. Washington DC: National Center for Education Statistics

Assessing for Literacy Tools

Rapid Test for Health Literacy - AHRQ



REALM-SF Score Sheet

Patient ID #: _____ Date: _____ Examiner Initials: _____

Behavior _____

Exercise _____

Menopause _____

Rectal _____

Antibiotics _____

Anemia _____

Jaundice _____

TOTAL SCORE _____

Score	Grade range
0	Third grade and below; will not be able to read most low-literacy materials; will need repeated oral instructions, materials composed primarily of illustrations, or audio or video tapes.
1-3	Fourth to sixth grade; will need low-literacy materials, may not be able to read prescription labels.
4-6	Seventh to eighth grade; will struggle with most patient education materials; will not be offended by low-literacy materials
7	High school; will be able to read most patient education materials

Informal Patient Assessments that may Help Identify Patient with Low Health Literacy – CHCS Fact Sheet



- “I forgot my reading glasses”
 - Frequently miss appts
 - Fail to complete registration forms
 - Unable to name medications or explain purpose or dosing
 - Identify pills by looking at them not reading label
 - Unable to give coherent, sequential medical history
 - Lack of follow through on tests/referrals
- Questions that may help HCP assess health literacy:
 - “A lot of people have trouble reading and remembering health information because it is difficult. Is this a problem for you?”
 - “What do you like to read? What do you rely on most to learn about health issues? Everyone has a unique source. TV? Radio? Internet? Friends and family?”

Impact of Low Health Literacy



- Medication errors
- Low rates of treatment adherence due to poor communication between providers and patients
- Reduced use of preventive services and more emergency room visits
- Poor management of chronic conditions
- Longer hospital stays
- More hospital visits (6% more)
- Less responsive to public health emergencies
- **4X higher health care cost**

CHCS Center for Health Care Strategies, Inc. FACT SHEET

Low Literacy in Patients With Diabetes



Study of 398 patients with diabetes: 83% of patients had previous diabetes education

- 25% could not determine what glucose meter readings were within normal range (80-120)
- 56% could not calculate total carbohydrate content in container of snack chips
- 59% could not accurately calculate insulin dose based on meal carbohydrate content and blood glucose level

Strategies for Addressing Low Health Literacy – Written Materials



- Simple wording, short messages, 4th to 6th grade reading level
- Information limited to key points; minimize disease statistics, anatomy, physiology
- Focus on key actions and desired behaviors
- Use pictures to help convey message rather than to decorate page
- Color-coding of tabular information to guide successful patient use of materials
- Encourage shared goal setting between patient and HCP

Low Literacy Diabetes Toolkit

Wolff K, Cavanaugh K, Rothman R. The Diabetes Literacy and Numeracy Education Toolkit (DLNET). *The Diabetes Educator* 2009; 35(2): 233-245. doi:10.1177/0145721709331945

PART 1

INTRO TO DIABETES


- Diabetes is a disease that causes you to have extra sugar in your blood (high sugar).

High sugar in the blood can cause you problems with:

- Poor vision or blindness
- Kidney disease
- Heart attacks or strokes
- Numbness, tingling or pain in your nerve endings
- Foot sores and foot pain
- Less blood flow
- Infections

- **But, control of your blood sugar can help to stop these problems!**

The Major Problems From Diabetes



1

PART 1



CARDI•OH
Ohio Cardiovascular and Diabetes Health Collaborative

Low Literacy Diabetes Toolkit

Wolff K, Cavanaugh K, Rothman R., et al

Practice One Serving Size

Use the label below:

What is the serving size? _____

How many carbohydrate grams are in each serving? _____

If you eat one serving, you will get _____ grams of carb.

Nutrition Facts	
Serving Size 2 crackers (14 g)	
Servings Per Container About 21	
Amount Per Serving	
Calories 60 Calories from Fat 15	
	% Daily Value*
Total Fat 1.5g	2%
Saturated Fat 0g	0%
Trans Fat 0g	
Cholesterol 0mg	0%
Sodium 70mg	3%
Total Carbohydrate 10g	3%
Dietary Fiber Less than 1g	3%
Sugars 0g	
Protein 2g	
Vitamin A 0%	• Vitamin C 0%
Calcium 0%	• Iron 2%
* Percent Daily Values are based on a diet of 2,000 calories. Your daily values may be higher or lower depending on your calorie needs:	
	Calories: 2,000 2,500
Total Fat	Less than 65g 80g
Sat Fat	Less than 20g 25g
Cholesterol	Less than 300mg 300mg
Sodium	Less than 2400mg 2400mg
Total Carbohydrate	300g 375g
Dietary Fiber	25g 30g

2 servings is _____ crackers

Add

_____ grams of carb from 1 serving

+ _____ grams of carb from 1 serving

= _____ grams of carb from 2 servings

1/2 serving is _____ crackers

_____ grams of carb from 1 serving

divided by 2

= _____ grams of carb from 1/2 serving

3

PART 9




INSULIN FOR SET DOSE PLUS CORRECTION

How Much Insulin Do I Take?

My long lasting insulin is: _____
 (Brand Name)

My short lasting insulin is: _____
 (Brand Name)



Before Breakfast:

1. Take _____ units of _____ (long lasting insulin at _____ am)
 2. Test blood sugar
 3. If blood sugar is below 70, eat 4 glucose tablets
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is over _____ take ____ units of _____ (short lasting insulin)

Before Lunch:

1. Test blood sugar
 2. If blood sugar is below 70, eat 4 glucose tablets
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is over _____ take ____ units of _____ (short lasting insulin)

Insulin For Set Dose Plus Correction



Before Supper:

1. Take _____ units of _____ (long lasting insulin)
 2. Test blood sugar
 3. If blood sugar is below 70, eat 4 glucose tablets
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)



At Bedtime:

1. Take _____ units of _____ (long lasting insulin)
 2. Test blood sugar
 3. If blood sugar is below 70, eat 4 glucose tablets
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is ____ to ____ take ____ units of _____ (short lasting insulin)
- If blood sugar is over _____ take ____ units of _____ (short lasting insulin)

Use Teach-Back Method

<https://www.ahrq.gov/health-literacy/improve/precautions/tool5.html>

AHRQ Health Literacy Universal Precautions Toolkit



- 1 – clinician gives a message or demonstrates a skill using plain language
- 2 – clinician asks patient to repeat the message using their own words or return demonstrate the skill to clinician
- 3 – any errors or misunderstandings are corrected by clinician and patient asked to repeat back the message or return demonstrate the skill

Avoid conveying disrespect to patient by taking responsibility for message:

“I want to make sure I’m explaining this clearly to you. Could you repeat back to me in your own words what I have explained to you?”

Teach Back

Please circle the foods that are part of a carbohydrate group.



Banana



Corn



Strawberries



Pepper



Pizza



Egg



Pancake



Broccoli



Spaghetti



Cheese



Peanut



Milk



Apple



Orange juice



Steak



Carrot



Tortilla



Pie



Ice Cream



Potato



Cake



Bread



Hamburger

Tips for Encounters

LEARN Model — clinical template for improved communication in cross-cultural patient-health care staff intervention



- L – Listen to patient’s perspective
- E – Explain and share one’s own perspective
- A – Acknowledge differences and similarities between the two perspectives
- R – Recommend treatment
- N – Negotiate mutually agreed upon plans

(Develop patient-provider relationship based on trust and respect)

Tips for Encounters with Patients with Low Literacy

CHCS Center for Health Care Strategies, Inc. FACT SHEET#5



- Create safe, respectful environment – greet warmly, make eye contact, get to know them, use cultural humility, earn trust, involve family/caretaker if possible, etc.
- Use simplified language

Instead of saying...	Say
hypertension	High blood pressure
pulmonary	Related to breathing
endocrinologist	Diabetes doctor
cardiac	Heart
HbA1c	Average blood sugar

Use Strengths-Based, Person-First Language When Talking to and About Patients



- Strengths-based language focuses on what people know and what they can do, rather than focusing on what they are not doing, making it more empowering
- Example – “Sam is taking less insulin than prescribed because of fear of hypoglycemia,” instead of “Sam is noncompliant with insulin taking.”
- Person-first language – “Sam has diabetes” instead of “Sam is diabetic.”

Low Literacy Resources

Scripps.org/diabetes

Giving Yourself Insulin



Insulin Pen Instructions



1. Wash your hands.



2. Attach pen needle.




3. Dial 2 units of insulin.




4. Waste 2 units of insulin.



5. Dial ordered insulin dose.



6. Select injection site and clean area.
Recommended sites:
A, B and C.



7. Inject insulin:
count to 10
before removing.



8. Place used needle
in sharps container.

Learn more about Scripps Whittier Diabetes Institute, visit scripps.org/diabetes or call 1-877-WHITTIER (944-8843).



Scripps.org/diabetes – Culturally Appropriate Materials



scripps.org/services/metabolic-conditions/diabetes/diabetes-professional-training/handouts

Updated information about COVID vaccines, visitor guidelines

[Doctors & Services](#) [Locations](#) [Patients](#)

[Home](#) > [Services](#) > [Diabetes Professional Training](#) > [Professional Handouts](#)

Diabetes Handouts for Providers

Multilingual diabetes handouts

[Expand All](#) | [Collapse All](#)

- [Arabic](#)
- [Chinese](#)
- [English](#)
- [Lao](#)
- [Somali](#)
- [Spanish](#)
- [Tagalog](#)
- [Vietnamese](#)

National Center for Farmworker Health – Bilingual Materials

Low Literacy English and Spanish

REGLAS PARA COMER SALUDABLEMENTE



1. Coma

Desayuno




Almuerzo



Cena

2. Si lo necesita,
pierda peso



3. Coma más fibra



4. Coma menos grasa.



Trate de no comer
comidas con mucha grasa!



RULES FOR HEALTHY EATING

5. Eat less sugar.



Try not to eat sweets.



6. Eat less salt.




Don't eat foods
with lots of salt.



7. Exercise every day.



8. Eat less meat.




Type 2 diabetes

In type 2 diabetes, the body can make some insulin, but not enough. Or, the insulin the body makes does not work right.

Type 2 diabetes often starts in adults, but children can have it too. It is more common in overweight people or if someone in the family has diabetes.

Type 2 diabetes is controlled by balancing when and how much you eat with:

- how active you are
- your weight, and
- the diabetes medicine you take



CARDI·OH
Ohio Cardiovascular and Diabetes Health Collaborative

learningaboutdiabetes.org

Low Literacy Resources

- <https://www.Learningaboutdiabetes.org>
- https://www.ncfh.org/health_education_resources.html
National Center for Farmworker Health – bilingual materials
- <https://www.cardi-oh.org/assets/qip/diabetes/patient-resources>
- <https://www.Scripps.org>

Free Program Through ADA diabetes.org

Project Power



Project Power for Adults

No-cost lifestyle change program.

Project Power is a no-cost type 2 diabetes lifestyle change program. Combining interactive lessons with a health coach, small support groups, and tools and resources, the program empowers you to reach your personal health goals.

[Sign up to Project Power for adults.](#)



Project Power for Youth

Healthy lifestyle program.


Project Power is a fun program with the aim to slow the trajectory of childhood obesity. This no-cost program, for kids ages 5–12, aims to help reduce its consequences. The program promotes making healthy food choices, increasing physical activity, and building family and peer support.

[Sign up to Project Power for youth.](#)

Use Technology to Assist With Low Numeracy Skills

- InPen
- Bigfoot Unity


Inpen compatible with Guardian CGM, Dexcom G5,6,7



Bigfoot Unity App



Black Cap for Long-Acting Insulin



White Cap for Rapid-Acting Insulin



Abbott FreeStyle Libre 2 Sensor



Bigfoot Blood Glucose Meter

References



- https://www.chcs.org/media/What_is_Health_Literacy.pdf
- Kutner M, Greenberg E, Jin Y, Paulsen C (2006). The Health Literacy of America's Adults: Results from the 2003 National Assessment of Adult Literacy (NCES 2006-483). U.S. Department of Education. Washington DC: National Center for Education Statistics
- Agency for Healthcare Research and Quality
<https://www.ahrq.gov/health-literacy/research/tools/index.html#rapid>
- https://www.huskyhealthct.org/providers/provider_postings/diabetes/DNT5.pdf Diabetes Numeracy Test
- Wolff K, Cavanaugh K, Rothman R. The Diabetes Literacy and Numeracy Education Toolkit (DLNET). The Diabetes Educator 2009; 35(2): 233-245. doi:10.1177/0145721709331945
- <https://www.ahrq.gov/health-literacy/improve/precautions/tool5.html>
- Ladha T, Zubairi M, Hunter A, Audcent T, Johnstone J. Cross-cultural communication: Tools for working with families and children. Paediatr Child Health. 2018 Feb;23(1):66-69. doi: 10.1093/pch/pxx126. Epub 2018 Feb 15. PMID: 29479280; PMCID: PMC5815116.
- National Standards for Diabetes Self-Management Education and Support. Diabetes Care 2022; Vol 45(2); 484-494. <https://doi.org/10.2337/dc21-2396>
- <https://www.Scripps.org/services/metabolic-conditions/diabetes/diabetes-professional-training/handouts>
- <https://www.Learningaboutdiabetes.org>
- <https://www.cardi-oh.org/assets/qip/diabetes/patient-resources>
- https://www.ncfh.org/health_education_resources.html



Thank you!

Questions/Discussion