

Mental Health and Chronic Conditions: Treating the Whole Patient to Improve Self-Care

Contributing authors on behalf of Team Best Practices:

Trygve Dolber, MD, Case Western Reserve University

Kate Gawlik, DNP, APRN-CNP, RN, The Ohio State University

Eileen Seeholzer, MD, MS, Case Western Reserve University

Bridget Lombard, DO, Ohio University

Shanna Stryker, MD, MPH, University of Cincinnati

Mental health and chronic conditions are frequently co-occurring. However simultaneously addressing these conditions can be complex and challenging.

Of the many mental health conditions, anxiety and depressive disorders are most common. The most effective treatment for anxiety and depression is often the combination of medication, counseling (such as cognitive behavioral therapy), and physical activity. Helping patients who have recurrent depression and anxiety create a long-term plan with these supports may be needed to sustain long-term cardiometabolic health and mental well-being.

The purpose of this document is to discuss the intersection between mental health and chronic conditions, specifically cardiometabolic disease. This document also discusses culture, stigma, and other barriers to receiving care for mental health conditions. Finally, evidence-based tools and interventions are reviewed to assist primary care providers in helping patients understand and address their conditions.



Importance of Identifying and Treating Mental Health Conditions in Patients With Cardiometabolic Disease

Psychological distress and mental illness are associated with increased risk of cardiovascular death, with rates of mental health disorders higher in patients with diabetes. Combined, cardiovascular disease (CVD) and mental illness are among the leading causes of morbidity and mortality worldwide.¹ Many mental health disorders and chronic diseases, specifically CVD and diabetes, have bidirectional relationships with shared biological, behavioral, psychological, and genetic mechanisms.¹⁻⁷ Mental health conditions impair self-management and likelihood of following a treatment plan, which can worsen cardiometabolic outcomes.¹

For minority populations, a lower frequency of these conditions is often reported. The lower rates are thought to be due to sociocultural factors, including stigma associated with mental illness, making these conditions less likely to be diagnosed or treated in patients who identify as part of a racial or ethnic minority. Having both cardiometabolic conditions and comorbid mental health condition(s) worsens clinical outcomes for both types of conditions.^{7,8}



- Individuals with common mental health conditions, such as major depressive disorder, have a 55% increased risk of developing CVD.¹
- Individuals with anxiety and posttraumatic stress disorder (PTSD) have a 27% increased risk of developing CVD.¹
- Individuals with PTSD have a two-fold increased risk of developing diabetes.²⁻⁶
- Individuals with diabetes, compared to those without diabetes, are two to three times more likely to have depression and are 20% more likely to have anxiety.⁷

Common Signs and Symptoms of Mental Health Disorders

Certain patterns of physical and behavioral symptoms can alert a provider that a patient may be struggling with a mental health condition. Unfortunately, the stigma and cultural norms about these conditions may lead patients to hide or feel ashamed of their symptoms and distress.

Signs and Symptoms of Anxiety and Depression

Physical

Anxiety or emotionality can exacerbate and cause physical symptoms to persist⁹⁻¹²

Somatization (in general)

- Any atypical complaint
- Symptoms out of proportion to pathology
- Limited ability to reassure
- Past acute event

Chronic Somatic Medical Conditions

- Irritable Bowel Syndrome
- Obesity¹³⁻²³
- Asthma exacerbations

Unexplained Somatic Symptoms

- Pelvic pain syndrome
- Chronic pain
- Cardiac symptoms including sinus tachycardia and recurrent chest pain if other causes have been ruled out
- Dizziness
- Headache
- Insomnia
- Fatigue
- Dyspepsia



Behavioral


- Abuse of and dependence on substances, including tobacco, cannabis, alcohol, or other substances²⁴
- Engagement in impulsive or risky behaviors that may manifest as:
high-risk sexual activity, frequent sexually transmitted infections, frequent or atypical legal trouble, failure to maintain healthy relationships, inability to maintain a job, or frequent car accidents
- Chronic anger and irritability
- Lack of engagement in recommended treatment plans
- Avoidance behaviors²⁵



The Impact of Stigma on Receiving Care

Stigma is defined as “the negative social attitude attached to a characteristic of an individual that may be regarded as a mental, physical, or social deficiency. A stigma implies social disapproval and can lead unfairly to discrimination against and exclusion of the individual.”²⁶ It is the fourth-highest ranked barrier to help-seeking.²⁷

Racial and ethnic minorities, youth, men, and those in the military and health professions report having a negative perception of healthcare as a result of stigmatization. How or whether one experiences stigma or internalizes shame because of stigmatization varies based on a wide number of intrinsic and extrinsic factors. Individual factors such as lived experience, race, culture, religion, gender, age, socioeconomic standing, and profession can impact the extent to which stigma affects health outcomes and personal wellness.²⁸



Common Reasons Cited for Not Seeking and Accepting Treatment for Mental Health Disorders

- Mistrust of health care providers due to longstanding inequalities in healthcare access and care (e.g., lack of culturally informed care, etc.)^{27, 29, 30, 31}
- Belief that mental health conditions violate the cultural value of being a hard-working person who must independently overcome and manage problems (e.g., males should be self-sufficient at managing mental health conditions)^{28, 31-34}
- Belief that antidepressant medications are addictive, change thinking processes or personality, or cause loss of control, and/or do physical harm³¹
- Hesitancy disclosing private thoughts and emotions to others^{31, 35}
- View that symptoms are a sign of insufficient faith, spiritual weakness, or bad karma from past misdeeds³¹
- Thought that a mental health condition will limit one’s competitiveness in obtaining or maintaining high-performing jobs³¹
- Negative attitudes toward mental health care and lack of perceived need for treatment^{30, 31}

Common Barriers to Making Changes to Address Health Issues

- Anhedonia or social withdrawal related to depression^{28,32}
- Disorganization related to anxiety (e.g., other stresses predominate, difficulty making decisions, etc.)³²
- Avoidance related to health anxiety (too overwhelming to confront disease)^{32, 36}
- Avoidance due to phobias³²
- Concern that a lifestyle change may deprive a person of unhealthy coping behaviors (e.g., stress eating, smoking) without offering an alternative way to manage stress and difficult feelings³²
- Distrust of provider or medical establishment in general; can be due to personal, familial, or historical trauma or negative experience³²
- Cost or financial barriers^{32, 36}
- Effort needed to find insurance coverage and “in network” clinicians³²



Managing Patients' Multiple Care Needs

Both clinicians and patients can be overwhelmed. Everything cannot be accomplished in one visit, but with the right approach, the initial conversation can end with empathy and hopefulness. The clinician can plan with a patient to discuss education and treatment options at subsequent visits.

Start by normalizing symptoms with the patient. Consider saying, *“Many patients struggle with feeling sad or down, overwhelmed, or worrying a lot. Is it possible that this is something you may be struggling with as well?”*

Ask for the patient’s thoughts about a particular mental health condition that may be impacting their health. Listening to their experience of symptoms and feelings about a mental health condition is the first step to overcoming barriers to accepting treatment for these conditions.

Patients can become more accepting of diagnosis and treatment of mental health conditions when they learn how mental health conditions and medical problems can interact. Patients should understand that if both types of conditions are not treated together, the patient’s health and quality of life may be much worse. For example, the clinician might say: *“We talked about how the mind’s automatic reaction to anxiety about blood sugar levels is to avoid checking those levels. But then, your diabetes gets worse, and your anxiety about diabetes becomes even more intense. To stop this cycle, we need to focus on treating anxiety and diabetes together, not just one or the other.”*

If short on time or overwhelmed, remember patient outcomes are improved when the clinician:

- Listens to a patient share their anxiety and/or encourages them to share with a loved one
- Educates the patient on how anxiety works and how treatment improves outcomes
- Helps patients to feel understood and to know that treatment improves health
- Uses the BATHE technique to communicate more effectively with patients³⁷

BATHE Technique³⁷

B = Background (e.g., Tell me more about this. What is going on in your life?)

A = Affect (e.g., How are you feeling about this?)

T = Trouble (e.g., What is the most difficult thing about this situation for you? What troubles you the most about this?)

H = Handling (e.g., How are you currently handling this problem/situation? How are you currently coping with this situation?)

E = Empathy (e.g., This must be hard for you. I can understand how this situation could be difficult for you.)

Strategies to Improve Trust in Patient-Provider Relationships

- Build patient trust and rapport early³⁸
- Make a positive first impression
- Ensure continuity of care (limiting the number of times a patient has to tell their story)
- Avoid judgmental language and behaviors
- Decrease anxiety by providing reassurance and empathy, encouraging patient questions, and avoiding medical jargon
- Demonstrate competence³⁹
 - Provide information about the medical condition, therapeutic regimen, and self-care
 - Provide directions and instructions for the therapeutic regimen
 - Ask for the patient's opinions and whether they understand
 - Partner with the patient in lieu of a paternalistic stance
 - Have familiarity with the patient's chart and condition
 - Clearly explain the rationale for tests and treatments
- Demonstrate warmth³⁹
 - Greet patient warmly and use the patient's name
 - Engage the patient in positive and/or social talk
 - Be receptive to the patient's emotions
 - Use non-verbal body language that portrays interest and concern including things like eye contact and tone of voice

For more information, visit Cardi-OH's expanded resources on [mental health](#), [reducing stigma](#) and [shared decision making](#).

Evidence-Based Interventions to Improve Mental Health and Cardiometabolic Outcomes

A patient-centered approach involves mutual decision making, inclusion of patient preferences and values, shared care planning, and patient participation in goal setting. All components are necessary for partnering with your patient to manage complex conditions.⁴⁰

Consider referring patients to a mental health care provider or counselor or providing one or more of the following interventions:

- **Psychoeducation** about the impact of worry and the connection between mental health symptoms and their medical disease(s). For example, “When distressing feelings or thoughts come up, the mind’s automatic reaction is to make them stop or get away from them. But if that distress is related to a disease, and if the reaction makes you avoid caring for it, the disease can progress.”
- **Methods of acceptance** that retrain the mind to navigate distress in a skillful way:
 - *Helpful thoughts*: “It does not matter if a thought is true or not, but is it helpful right now? If not, find a way to let it go and do what you need to do.”
 - *Relaxation techniques*: Diaphragmatic breathing, progressive muscle relaxation
 - *Curiosity practice*: Open up to distress and explore how it feels without judgment. The free app, Mindfulness Coach, can help the patient learn.⁴¹
- **Cognitive Behavioral Therapy**
- **Medications for depression or anxiety**
- **Care Coordination / Team Based Care Models** (e.g., Collaborative Chronic Care Model)⁴²⁻⁴⁶
- **Self-help tools** including evidence-based stress reduction apps, websites, and books. Durable and positive effects on mindfulness, general psychiatric symptoms, and several aspects of quality of life at low costs are possible with smartphone apps.
- **Lifestyle changes** that can improve symptoms (e.g., exercise, sleep hygiene, healthy eating, reducing substance use)

The most effective treatment for anxiety and depression is often the combination of medication, counseling (such as cognitive behavioral therapy), and physical activity together. Helping patients who have recurrent depression and anxiety create a long-term plan with these supports may be needed to sustain long-term cardiometabolic health and mental well-being.

References:

1. De Hert M, Detraux J, Vancampfort D. The intriguing relationship between coronary heart disease and mental disorders. *Dialogues Clin Neurosci*. 2018;20(1):31-40. doi: 10.31887/DCNS.2018.20.1/mdehert.
2. Scherrer JF, Salas J, Lustman PJ, et al. The role of obesity in the association between posttraumatic stress disorder and incident diabetes. *JAMA Psychiatry*. 2018;75(11):1189-1198. doi:10.1001/jamapsychiatry.2018.2028.
3. Goldstein RB, Smith SM, Chou SP, et al. The epidemiology of DSM-5 posttraumatic stress disorder in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions-III. *Soc Psychiatry Psychiatr Epidemiol*. 2016;51(8):1137-1148. doi:10.1007/s00127-016-1208-5.
4. Rosenbaum S, Stubbs B, Ward PB, et al. The prevalence and risk of metabolic syndrome and its components among people with posttraumatic stress disorder: a systematic review and meta-analysis. *Metabolism*. 2015;64(8):926-933. doi:10.1016/j.metabol.2015.04.009.
5. Miller-Archie SA, Jordan HT, Ruff RR, et al. Posttraumatic stress disorder and new-onset diabetes among adult survivors of the World Trade Center disaster. *Prev Med*. 2014;66:34-38. doi: 10.1016/j.jypmed.2014.05.016.
6. Boyko EJ, Jacobson IG, Smith B, et al. Risk of diabetes in U.S. military service members in relation to combat deployment and mental health. *Diabetes Care*. 2010;33(8):1771-1777. doi:10.2337/dc10-0296.
7. Centers for Disease Control and Prevention. Diabetes and Mental Health. <https://www.cdc.gov/diabetes/managing/mental-health.html>. Reviewed May 7, 2021. Accessed September 6, 2021.
8. Centers for Disease Control and Prevention. Diabetes Report Card 2019. www.cdc.gov/diabetes/library/reports/congress.html. Reviewed January 4, 2021. Accessed September 5, 2021.
9. Trivedi MH. The link between depression and physical symptoms. *Prim Care Companion J Clin Psychiatry*. 2004;6(Suppl 1):12-16. PMID: PMC486942.
10. Bagayogo IP, Interian A, Escobar JI. Transcultural aspects of somatic symptoms in the context of depressive disorders. *Adv Psychosom Med*. 2013;33:64-74. doi: 10.1159/000350057.
11. Kessler D, Bennewith O, Lewis G, Sharp D. Detection of depression and anxiety in primary care: follow up study. *BMJ*. 2002; 325 :1016. doi:10.1136/bmj.325.7371.1016.
12. Johnson EO, Roth T, Breslau N. The association of insomnia with anxiety disorders and depression: exploration of the direction of risk. *J Psychiatr Res*. 2006;40(8):700-708. doi:10.1016/j.jpsychires.2006.07.008.
13. Mitchell KS, Aiello AE, Galea S, et al. PTSD and obesity in the Detroit neighborhood health study. *Gen Hosp Psychiatry*. 2013;35(6):671-673. doi:10.1016/j.genhosppsych.2013.07.015.
14. de Zwaan, M. Binge eating disorder and obesity. *Int J Obes* 25, S51–S55 (2001). <https://doi.org/10.1038/sj.ijo.0801699>.
15. Gelenberg AJ. Psychiatric and somatic markers of anxiety: identification and pharmacologic treatment. *Prim Care Companion J Clin Psychiatry*. 2000;2(2):49-54. doi:10.4088/pcc.v02n0204.
16. Garipey G, Nitka D, Schmitz N. The association between obesity and anxiety disorders in the population: a systematic review and meta-analysis. *Int J Obes (Lond)*. 2010;34(3):407-419. doi:10.1038/ijo.2009.252.
17. Del Giacco SR, Cappai A, Gambula L, et al. The asthma-anxiety connection. *Respiratory Medicine*. 120 (2016): 44-53. <https://doi.org/10.1016/j.rmed.2016.09.014>.
18. Bytzer P, Talley NJ. Dyspepsia. *Ann Intern Med*. 2001 May 1;134(9 Pt 2):815-22. doi: 10.7326/0003-4819-134-9_part_2-200105011-00004.
19. Pontari MA, Ruggieri MR. Mechanisms in prostatitis/chronic pelvic pain syndrome. *J Urol*. 2004;172(3):839-845. doi:10.1097/01.ju.0000136002.76898.04.
20. Latthe P, Mignini L, Gray R, et al. Factors predisposing women to chronic pelvic pain: systematic review. *BMJ*. 2006;332(7544):749-755. doi:10.1136/bmj.38748.697465.55.
21. Leeuw M, Goossens ME, Linton SJ, et al. The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J Behav Med*. 2007;30(1):77-94. doi:10.1007/s10865-006-9085-0.
22. Shabtaie SA, Witt CM, Asirvatham SJ. Natural history and clinical outcomes of inappropriate sinus tachycardia. *J Cardiovasc Electrophysiol*. 2020;31(1):137-143. doi:10.1111/jce.14288.
23. Eifert GH, Zvolensky MJ, Lejuez CW. Heart-focused anxiety and chest pain: a conceptual and clinical review. *Clinical Psychology: Science and Practice*, 7(4), 403–417. <https://doi.org/10.1093/clipsy.7.4.403>.
24. Robinson J, Sareen J, Cox BJ, Bolton J. Self-medication of anxiety disorders with alcohol and drugs: results from a nationally representative sample. *J Anxiety Disord*. 2009;23(1):38-45. doi:10.1016/j.janxdis.2008.03.013.
25. Kryptos AM, Efftig M, Kindt M, Beckers T. Avoidance learning: a review of theoretical models and recent developments. *Front Behav Neurosci*. 2015;9:189. doi:10.3389/fnbeh.2015.00189.
26. American Psychological Association. APA Dictionary of Psychology. <https://dictionary.apa.org/stigma>. Accessed April 7, 2022.

27. Clement S, Schauman O, Graham T, et al. What is the impact of mental health-related stigma on help-seeking? A systematic review of quantitative and qualitative studies. *Psychol Med*. 2015 Jan;45(1):11-27. doi: 10.1017/S0033291714000129.
28. Goetter EM, Frumkin MR, Palitz SA, et al. Barriers to mental health treatment among individuals with social anxiety disorder and generalized anxiety disorder. *Psychol Serv*. 2020 Feb;17(1):5-12. doi: 10.1037/ser0000254.
29. Henderson C, Noblett J, Parke H, et al. Mental health-related stigma in health care and mental health-care settings. *Lancet Psychiatry*. 2014;1(6):467-482. doi:10.1016/S2215-0366(14)00023-6.
30. Harris EC, Barraclough B. Excess mortality of mental disorder. *Br J Psychiatry*. 1998;173:11-53. doi:10.1192/bjp.173.1.11.
31. Yang LH, Thornicroft G, Alvarado R, et al. Recent advances in cross-cultural measurement in psychiatric epidemiology: utilizing D23 'what matters most' to identify culture-specific aspects of stigma. *Int J Epidemiol*. 2014 Apr;43(2):494-510. doi: 10.1093/ije/dyu039.
32. Hines-Martin V, Malone M, Kim S, Brown-Piper A. Barriers to mental health care access in an African American population. *Issues Ment Health Nurs*. 2003;24(3):237-256. doi:10.1080/01612840305281.
33. Milner A, Law PCF, Mann C, et al. A smart-phone intervention to address mental health stigma in the construction industry: a two-arm randomised controlled trial. *SSM Popul Health*. 2017 Dec 31;4:164-168. doi: 10.1016/j.ssmph.2017.12.007.
34. Latalova K, Kamaradova D, Prasko J. Perspectives on perceived stigma and self-stigma in adult male patients with depression. *Neuropsychiatr Dis Treat*. 2014 Jul 29;10:1399-405. doi: 10.2147/NDT.S54081.
35. Kaushik A, Kostaki E, Kyriakopoulos M. The stigma of mental illness in children and adolescents: a systematic review. *Psychiatry Res*. 2016 Sep 30;243:469-94. doi: 10.1016/j.psychres.2016.04.042.
36. Lavingia R, Jones K, Asghar-Ali AA. A systematic review of barriers faced by older adults in seeking and accessing mental health care. *J Psychiatr Pract*. 2020 Sep;26(5):367-382. doi: 10.1097/PRA.0000000000000491.
37. Thomas C, Cramer H, Jackson S, et al. Acceptability of the BATHE technique amongst GPs and frequently attending patients in primary care: a nested qualitative study. *BMC Fam Pract*. 2019;20(1):121. doi:10.1186/s12875-019-1011-y.
38. Dang BN, Westbrook RA, Njue SM, Giordano TP. Building trust and rapport early in the new doctor-patient relationship: a longitudinal qualitative study. *BMC Med Educ*. 2017;17(1):32. doi:10.1186/s12909-017-0868-5
39. Howe LC, Leibowitz KA, Crum AJ. When your doctor "gets it" and "gets you": the critical role of competence and warmth in the patient-provider interaction. *Front Psychiatry*. 2019;10:475. doi:10.3389/fpsy.2019.00475.
40. Reiss-Brennan B, Brunisholz KD, Dredge C, et al. Association of integrated team-based care with health care quality, utilization, and cost. *JAMA*. 2016 Aug 23-30;316(8):826-34. doi: 10.1001/jama.2016.11232.
41. van Emmerik AAP, Berings F, Lancee J. Efficacy of a mindfulness-based mobile application: a randomized waiting-list controlled trial. *Mindfulness (N Y)*. 2018;9(1):187-198. doi:10.1007/s12671-017-0761-7.
42. Berry LL, Rock BL, Smith Houskamp B, et al. Care coordination for patients with complex health profiles in inpatient and outpatient settings. *Mayo Clin Proc*. 2013 Feb;88(2):184-94. doi: 10.1016/j.mayocp.2012.10.016.
43. Owens-Gary MD, Zhang X, Jawanda S, et al. The importance of addressing depression and diabetes distress in adults with type 2 diabetes. *J Gen Intern Med*. 2019;34(2):320-324. <https://doi.org/10.1007/s11606-018-4705-2>.
44. Germack HD, Leung L, Zhao X, et al. Association of team-based care and continuity of care with hospitalizations for veterans with comorbid mental and physical health conditions. *J Gen Intern Med*. 2022;37(1):40-48. doi: 10.1007/s11606-021-06884-5.
45. Miller CJ, Griffith KN, Stolzmann K, et al. An economic analysis of the implementation of team-based collaborative care in outpatient general mental health clinics. *Med Care*. 2020 Oct;58(10):874-880. doi: 10.1097/MLR.0000000000001372.
46. Reiss-Brennan B. Mental health integration: normalizing team care. *J Prim Care Community Health*. 2014 Jan 1;5(1):55-60. doi: 10.1177/2150131913508983.

Partners



In partnership with:



The Ohio Cardiovascular & Diabetes Health Collaborative is funded by the Ohio Department of Medicaid and administered by the Ohio Colleges of Medicine Government Resource Center. The views expressed in this document are solely those of the authors and do not represent the views of the state of Ohio or federal Medicaid programs.