



# Addressing Health Literacy and Numeracy: A Focus on Nutrition

## Contributing authors on behalf of Team Best Practices:

Mary Beth Vonder Meulen, RN, University of Cincinnati

Rosellen Roche, MD, PhD, Ohio University

Kate Gawlik, DNP, APRN-CNP, The Ohio State University

Eileen Seeholzer, MD, MS, Case Western Reserve University

In the U.S., 54% of adults read below a sixth-grade level, and only 12% of U.S. adults have the health literacy skills needed to manage the demands of our complex health care system.<sup>1</sup>

Stress or illness can further compromise these individuals' ability to absorb and use health information.<sup>2</sup> Patients with low health literacy and numeracy are more likely to have poor health outcomes, lower medication adherence, delays in seeking care, increased hospitalizations, higher mortality, and higher health care costs than those with higher health literacy.<sup>3,4</sup>

A person's ability to follow dietary recommendations (e.g., reading food labels, calculating nutrition information, establishing schedules for eating around medications, etc.) is particularly crucial for managing chronic diseases such as diabetes and cardiovascular disease, given the addition of pharmacotherapies. Patients with higher nutrition literacy and numeracy skills are more likely to interpret food labels correctly and have better health outcomes.<sup>7</sup>

This document identifies screening tools for health literacy and numeracy, provides guidance on how to use nutrition food labels in patient education, and offers strategies to improve patient comprehension of medical information.

## HEALTH LITERACY

Health literacy is the degree to which individuals can find, understand, and use information and services to inform health-related decisions and actions for themselves and others.<sup>5</sup>

**Examples:** Finding and accessing care, processing verbal medical instructions, and making informed decisions).

## HEALTH NUMERACY

Health numeracy refers to the ability to understand clinical and public health data subjectively and/or apply it objectively.<sup>6</sup>

**Examples:** Calculating portion sizes or nutrition information

Patients with low health literacy and numeracy are likely to have difficulty understanding and processing medical information needed for their own care or the care of a loved one.

## Health Literacy and Numeracy Screening Tools

Health literacy and numeracy assessments are foundational components for achieving excellent patient outcomes. While screening for health literacy and numeracy has not yet been shown to improve health outcomes, consensus statements support universal health literacy and numeracy precautions. This means “providing understandable and accessible information to all patients, regardless of their literacy or education levels.”<sup>8</sup> For patients needing to adopt healthy diet changes and increase activity long-term, health literacy and numeracy are needed to make and monitor choices.

Using the following resources, primary care teams can choose a method to assess their patient’s health literacy and numeracy and adopt recommended strategies to tailor conversations.

**Agency for Healthcare Research and Quality (AHRQ) Health Literacy Universal Precautions Toolkit:** A wide range of resources is available in this toolkit, including assessment tools for individuals and practices, tools to improve patient engagement and education, and training for the teach-back method.<sup>9</sup>

[ahrq.gov/health-literacy/improve/precautions/index.html](http://ahrq.gov/health-literacy/improve/precautions/index.html)

**Health Literacy Tool Shed:** This collection of assessment tools, funded by the National Institute of Health’s National Library of Medicine, can be narrowed down to specific diseases such as diabetes. For example, the Diabetes Numeracy Test evaluates a wide range of numeracy skills used by patients with diabetes, especially the ability to use the Nutrition Facts Panel (NFP) label.<sup>10</sup>

[healthliteracy.bu.edu/specific\\_context=6](http://healthliteracy.bu.edu/specific_context=6)

**Newest Vital Sign:** This validated test developed by Pfizer uses a nutrition label and six questions to quickly assess health literacy and numeracy in English and Spanish.<sup>8,11</sup> The test only takes three minutes to administer and requires both reading and math skills to complete. A score less than four indicates patients are at high risk for low health literacy.

[cdn.pfizer.com/pfizercom/health/nvs\\_flipbook\\_english\\_final.pdf](http://cdn.pfizer.com/pfizercom/health/nvs_flipbook_english_final.pdf)

**Single Item Literacy Screener:** This validated tool can be used in the clinic to identify patients who may need help reading health materials. The team can easily collect the information in about 15 seconds while taking vital signs.<sup>12</sup>

### Single Item Literacy Screener

“How often do you need to have someone help you when you read instructions, pamphlets, or other written material from your doctor or pharmacy?”

1 – Never	2 – Rarely	3 – Sometimes
4 – Often	5 – Always	

Scoring:  $\geq$  indicates some difficulty reading, printed health related material (i.e., positive screen)

## Understanding Nutrition Food Labels

It is important for primary care teams to assess a patient's health literacy/numeracy and use methods and tools to help these patients better manage their health conditions and participate in collaborative decision making about their care. Nutrition education using simplified teaching methods that emphasize understanding food labels has been shown to improve blood pressure and blood glucose levels.<sup>13,14</sup>

### Nutrition Facts Panel

The Nutrition Facts Panel (NFP), developed by the U.S. Food and Drug Administration (FDA) in 1990 and revised in 2016, was intended to provide information to the public needed to follow dietary recommendations and make healthier food choices.<sup>15</sup> However, the NFP has been criticized as too complex for many consumers, particularly those with low health literacy and numeracy.<sup>15-18</sup>

Recognizing this, the FDA now provides educational information about the new food label for teachers, parents, and other consumers.<sup>15</sup> Most have links to PDFs that can be used in the office, including information about what percent Daily Value (%DV) is considered high and low. However, many of these resources assume a proficient reading level and access to a computer or smartphone with sufficient data access. Primary care providers can use the NFP in addition to a visual dietary guide like [MyPlate](#) or the [Cardi-OH DASH Diet Plate](#) to assist patients with limited health literacy and numeracy.

Patient understanding can be reinforced through guidance, repeat visits, and practice, focusing on a few prioritized areas on the label at any given appointment. These areas can then be correlated to better choices on the plate. For example, grams of carbohydrate can be discussed at an office visit with labels and teach-back for a basic goal for meals (i.e., 30-45 grams of carbohydrate per meal). This iterative and step-wise approach can help patients to improve and adapt choices and dietary patterns practically, moving toward healthier eating patterns over time.

Nutrition Facts	
4 servings per container	
Serving size 1 cup (227g)	
Amount per serving	
<b>Calories 280</b>	
% Daily Value*	
<b>Total Fat</b> 9g	12%
Saturated Fat 4.5g	23%
Trans Fat 0g	
<b>Cholesterol</b> 35mg	12%
<b>Sodium</b> 850mg	37%
<b>Total Carbohydrate</b> 34g	12%
Dietary Fiber 4g	14%
Total Sugars 6g	
Includes 0g Added Sugars	0%
<b>Protein</b> 15g	
Vitamin D 0mcg	0%
Calcium 320mg	25%
Iron 1.6mg	8%
Potassium 510mg	10%

\* The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 2,000 calories a day is used for general nutrition advice.

■ Servings Per Container    ■ % Daily Value (%DV)  
■ Serving Size    ■ Calories

Use %DV to determine if a food serving is high ( $\geq 20\%$  DV) or low ( $\leq 5\%$  DV) in an individual nutrient.

## Front of Package Label

The Front of Package (FOP) label, also known as the Facts Up Front label, is voluntarily adopted by food manufacturers and has been used worldwide since 2011. The FOP's simple, easy-to-use format highlights key information from the NFP label about calories, saturated fat, sodium, and sugar – nutrients recommended to be limited in the Dietary Guidelines for Americans.<sup>19</sup> The four nutrient facts are always presented together as a consistent set.<sup>20</sup>

### PER 1/2 CUP



Adapted from Facts Up Front<sup>21</sup>

Primary care teams may use the FOP label to provide patients with daily goals and limits that require only basic arithmetic to understand, giving patients with low literacy and numeracy the ability to make healthier food choices. The FOP label can be quickly reviewed as part of a clinical visit, giving the patient a useful tool to use until they can connect with a registered dietitian or **Diabetes Self-Management Education and Support (DSMES)** provider or program.

For more information, access Cardi-OH's expanded resources on **health literacy, addressing disparities** in cardiovascular disease and diabetes, and **personalizing care** to optimize outcomes.

## Strategies for Primary Care Teams to Improve Patient Comprehension of Medical Information

Accessible and effective health literacy tools and information equip patients to participate in shared decision making, select recommended care plans, and adhere to health and lifestyle choices. When patients are identified as having low health literacy and/or numeracy, primary care teams can use targeted interventions to improve their patients' understanding and promote improved health literacy and numeracy. Some evidence-based strategies include:<sup>22-30</sup>

### Communicating with Patients:

- Use jargon-free language when communicating with patients.
- Focus on key messages and consistently repeat them in written, verbal, and visual approaches
- Ask patients to repeat back what they learned using the teach-back method.
- Adapt or design written materials for the sixth-grade reading level or lower.
- Adopt technology and eHealth interventions in patient education (i.e., videos and interactive self-help tools) for use during visits and in medical waiting or counseling areas. At home, patients can search for educational videos about “food labels” on **YouTube**, offered by official government or educational institutions, such as the **FDA** or **American Diabetes Association**.
- Encourage regular follow-up visits to improve understanding and answer questions.
- Use interpreter services if a language barrier exists.
- Engage family members, caregivers, and support persons in patient education sessions and equip them with appropriate resources.
- Refer to one-on-one educational sessions with a specialist (i.e., diabetes educator, pharmacist, registered dietitian).

### Enhancing Provider Skills and Teamwork:

- Have primary care teams complete health literacy training (e.g. the basic use of the NFP and FOP).
- Implement a team-based approach for screening and teaching for low health literacy and numeracy.

## References

1. U.S. Department of Education, National Center for Education Statistics. Highlights of PIAAC 2017 U.S. Results (NCES 2020-777). [https://nces.ed.gov/surveys/piaac/current\\_results.asp](https://nces.ed.gov/surveys/piaac/current_results.asp). Accessed July 25, 2022.
2. Agency for Healthcare Research and Quality. Health Literacy. <https://www.ahrq.gov/health-literacy/index.html>. Accessed July 25, 2022.
3. Bailey SC, Brega AG, Crutchfield TM, et al. Update on health literacy and diabetes. *Diabetes Educ.* 2014;40(5):581–604. doi:10.1177/0145721714540220.
4. Altin SV, Finke I, Kautz-Freimuth S, Stock S. The evolution of health literacy assessment tools: a systematic review. *BMC Public Health.* 2014;14:1207. doi:10.1186/1471-2458-14-1207.
5. Centers for Disease Control and Prevention. What is Health Literacy? <https://www.cdc.gov/healthliteracy/learn/index.html>. Reviewed February 2, 2022. Accessed October 3, 2022.
6. Centers for Disease Control and Prevention. Numeracy. <https://www.cdc.gov/healthliteracy/researchevaluate/numeracy.html>. Reviewed March 18, 2022. Accessed October 3, 2022.
7. Centers for Disease Control and Prevention. Food Literacy. <https://www.cdc.gov/healthliteracy/researchevaluate/food-literacy.html>. Updated May 21, 2021. Accessed December 27, 2021.
8. Hersh L, Salzman B, Snyderman D. Health literacy in primary care practice. *Am Fam Physician.* 2015 Jul 15;92(2):118-24. PMID:26176370.
9. Agency for Healthcare Research and Quality. Health Literacy Universal Precautions Toolkit, 2nd Edition. <https://www.ahrq.gov/health-literacy/improve/precautions/toolkit.html>. Reviewed September 2020. Accessed October 3, 2022.
10. National Institute of Health. Health Literacy Toolshed. <http://healthliteracy.bu.edu/>. Updated January 3, 2022. Accessed January 3, 2022.
11. Weiss B, Mays M, Martz W, et al. Quick assessment of literacy in primary care: the newest vital sign. *Ann Fam Med.* 2005; 3(6) 514-522. doi:10.1370/afm.405.
12. Morris NS, MacLean CD, Chew LD, Littenberg B. The Single Item Literacy Screener: evaluation of a brief instrument to identify limited reading ability. *BMC Fam Pract.* 2006;7:21. Published 2006 Mar 24. doi:10.1186/1471-2296-7-21.
13. Dang YH, Patel-Shori NM, Barros M, Yu D. Good literacy to enhance response in diabetes mellitus (GLITTER-DM). *Innov Pharm.* 2020 Jul 31;11(3):10.24926/iip.v11i3.2406. doi:10.24926/iip.v11i3.2406.
14. Warren-Findlow J, Coffman MJ, Thomas EV, Krinner LM. ECHO: A pilot health literacy intervention to improve hypertension self-care. *Health Lit Res Pract.* 2019 Dec 5;3(4):e259-e267. doi:10.3928/24748307-20191028-01.
15. Persoskie A, Hennessy E, Nelson WL. US consumers' understanding of nutrition labels in 2013: the importance of health literacy. *Prev Chronic Dis.* 2017 Sep 28;14:E86. doi:10.5888/pcd14.170066.
16. Rothman RL, Housam R, Weiss H, et al. Patient understanding of food labels: the role of literacy and numeracy. *Am J Prev Med.* 2006; 31(5): 391–398. doi:10.1016/j.amepre.2006.07.025.
17. Cha E, Kim KH, Lerner HM, et al. Health literacy, self-efficacy, food label use, and diet in young adults. *Am J Health Behav.* 2014; 38(3):331–339. doi:10.5993/AJHB.38.3.2.
18. Gibbs HD, Bonenberger H, Hull HR, et al. Validity of an updated nutrition literacy assessment instrument with the new nutrition facts panel. *Int J Food Sci Nutr.* 2020;71(1):116-121. doi:10.1080/09637486.2019.1606167.
19. U.S. Food and Drug Administration. Food Labeling & Nutrition. <https://www.fda.gov/food/food-labeling-nutrition>. Updated November 1, 2021. Accessed December 27, 2021.
20. Crockett RA, King SE, Marteau TM, et al. Nutritional labelling for healthier food or non-alcoholic drink purchasing and consumption. *Cochrane Database Syst Rev.* 2018; 2(2):CD009315. doi:10.1002/14651858.CD009315.pub2.
21. Facts Up Front. <http://www.factsupfront.org>. Accessed December 27, 2021.
22. Janz NK, Li Y, Zikmund-Fisher BJ, et al. The impact of doctor-patient communication on patients' perceptions of their risk of breast cancer recurrence. *Breast Cancer Res Treat.* 2017;161(3):525-535. doi:10.1007/s10549-016-4076-5.
23. Berkman ND, Sheridan SL, Donahue KE, et al. AHRQ Evidence Report/Technology Assessment Number 199, Health Literacy Interventions and Outcomes: an Updated Systematic Review. <https://www.ahrq.gov/downloads/pub/evidence/pdf/literacy/literacyup.pdf>. Published March 2011. Accessed July 25, 2022.
24. Jacobs RJ, Lou JQ, Ownby RL, Caballero J. A systematic review of eHealth interventions to improve health literacy. *Health Informatics J.* 2016;22(2):81-98. doi:10.1177/1460458214534092.
25. Pignone M, DeWalt DA, Sheridan S, et al. Interventions to improve health outcomes for patients with low literacy. A systematic review. *J Gen Intern Med.* 2005;20(2):185-92. doi:10.1111/j.1525-1497.2005.40208.x.
26. Smith SG, Kobayashi LC, Wolf MS, et al. The associations between objective numeracy and colorectal cancer screening knowledge, attitudes and defensive processing in a deprived community sample. *J Health Psychol.* 2016;21(8):1665-1675. doi:10.1177/1359105314560919.
27. van Weert JCM, Alblas MC, van Dijk L, Jansen J. Preference for and understanding of graphs presenting health risk information. The role of age, health literacy, numeracy and graph literacy. *Patient Educ Couns.* 2021;104(1):109-117. doi:10.1016/j.pec.2020.06.031.
28. Ha Dinh TT, Bonner A, Clark R, et al. The effectiveness of the teach-back method on adherence and self-management in health education for people with chronic disease: a systematic review. *JBIC Database System Rev Implement Rep.* 2016;14(1):210-247. doi:10.11124/jbisrir-2016-2296.
29. Black RL, Duval C. Diabetes discharge planning and transitions of care: a focused review. *Curr Diabetes Rev.* 2019;15(2):111-117. doi:10.2174/1573399814666180711120830.
30. Caruso R, Magon A, Baroni I, et al. Health literacy in type 2 diabetes patients: a systematic review of systematic reviews. *Acta Diabetol.* 2018;55(1):1-12. doi:10.1007/s00592-017-1071-1.cc.



---

## Partners



*In partnership with*



---

The Ohio Cardiovascular & Diabetes Health Collaborative is funded by the Ohio Department of Medicaid and administered by the Ohio Colleges of Medicine Government Resource Center. The views expressed in this document are solely those of the authors and do not represent the views of the state of Ohio or federal Medicaid programs.