

UPDATED MAY 2025 — CAPSULE 2

Home Blood Pressure (BP) Monitoring: Practical Instructions for Patients

CONTRIBUTING AUTHORS: Shari Bolen, MD, MPH; Shireen Khoury, MD, MPH; and Jackson T. Wright Jr., MD, PhD, on behalf of Team Best Practices

How many blood pressure (BP) readings should a patient take at home to inform a hypertension treatment decision?

Home BP monitoring is an important step in the evaluation and management of hypertension. Clinicians can facilitate more meaningful home BP readings by educating patients on how to measure their BP accurately.

Cardi-OH has developed **practical instructions** for accurate home BP monitoring to share with patients. Consider adapting these tips for use in clinical after-visit instructions.



When Should You Measure Your Blood Pressure?

In the 3 to 5 days before your next visit, take your blood pressure according to the schedule below and record all blood pressure readings in a log.¹ If your BP medication is changed, check weekly starting two weeks after the change and continuing until the next visit.²



Each day, take:

Two blood pressure readings in the morning, before taking your medications and two blood pressure readings in the evening, before going to bed.³

How to Take Your Blood Pressure at Home:

1. Do NOT smoke, exercise, or drink caffeine in the 30 minutes before checking your blood pressure.
2. Sit in a chair with your back supported and feet flat on the floor.
3. Rest your arm on a table at heart level.
4. Sit at rest for at least 5 minutes before taking the first blood pressure reading.
5. Do NOT talk during the 5 minute rest period or while taking your blood pressure.
6. Wait 1 minute before taking the second reading each time (morning and evening).
7. Write down the two readings along with the date and time.

Remember!

Bring both your log and home blood pressure monitor to each clinic visit to be checked against your doctor's office machines.

References

1. Bello NA, Schwartz JE, Kronish IM, et al. Number of measurements needed to obtain a reliable estimate of home blood pressure: results from the improving the detection of hypertension study. *J Am Heart Assoc.* 2018;7(20):e008658. doi:10.1161/JAHA.118.008658.
2. Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension.* 2018;71(6):1269-1324. doi:10.1161/HYP.0000000000000066.
3. Muntner P, Shimbo D, Carey RM, et al. Measurement of blood pressure in humans: a scientific statement from the American Heart Association. *Hypertension.* 2019;73(5):e35-e66. doi:10.1161/HYP.0000000000000087.

The Ohio Cardiovascular and Diabetes Health Collaborative is funded by the Ohio Department of Medicaid and administered by the Ohio Colleges of Medicine Government Resource Center. The views expressed in this document are solely those of the authors and do not represent the views of the state of Ohio or federal Medicaid programs.

For more information head to **Cardi-OH.org**.

© 2020 Cardi-OH