

JANUARY 2023 – CAPSULE 29

Using SMART Goals to Promote a Healthy Body Weight

CONTRIBUTING AUTHORS: James J. Werner, PhD, MSSA, Case Western Reserve University; Karen Bailey, MS, RD, LD, CDCES, Ohio University; Colin Crowe, MD, Case Western Reserve University, on behalf of Team Best Practices

When patients express a desire to lose weight or improve overall health, clinicians can guide patients in aligning their goals with each of the **SMART** (Specific, Measurable, Achievable, Relevant, Time-Based) goal criteria. Weight loss can be achieved and maintained by making lifestyle changes that benefit both weight and overall health.

- Ask the patient about their reasons for wanting to lose weight and express empathy by reflecting on their responses. Be aware that the patient may be sensitive to judgment or negativity, as many have experienced weight-based stigma in health care settings.¹
- Although it is common for patients to hope to achieve significant weight loss, advise that even a modest loss of 5 to 10% is likely to produce health benefits, including improvements in blood pressure, cholesterol, and glucose.²⁻⁴
- Explain SMART goal setting and its value. Invite the patient to choose one lifestyle behavior to implement. Evidence-based recommendations include:⁵⁻⁸
 - Eat a protein-rich breakfast as the biggest daily meal.
 - Drink only water, black coffee, or plain tea.
 - Stop all food intake at 7 p.m.
 - Exercise for 30 minutes daily.
- Inquire about anticipated obstacles to implementation. Using motivational interviewing techniques, elicit the patient's ideas about overcoming obstacles and assist with contingency planning.
- Plan for follow-up in four weeks to discuss SMART goal progress.
- Optional: Assist the patient in setting weekly and monthly weight loss goals. Weight loss should not exceed 2 pounds per week. Inquire about support for weight loss efforts. If resources are available, offer support from a community health worker or a member of the practice staff.

Specific: I plan to drink no other beverages besides water and black coffee on at least four of the next seven days.

Measurable: Each day that I achieve my goal, I will circle the date on my calendar before going to bed.

Achievable: This goal seems realistic and I believe that I can achieve it.

Relevant: Achieving this goal can help me lose weight and improve my health, which are important to me.

Time-Based: I plan to work on this goal for the next seven days.

For more information, access Cardi-OH's patient handout on **SMART goals** and expanded resources on **motivational interviewing** and **healthy lifestyle changes**.

References

1. Phelan SM, Burgess DJ, Yeazel MW, et al. Impact of weight bias and stigma on quality of care and outcomes for patients with obesity. *Obes Rev*. 2015;16(4):319-326. doi:10.1111/obr.12266.
2. Foster GD, Wadden TA, Vogt RA, et al. What is a reasonable weight loss? Patients' expectations and evaluations of obesity treatment outcomes. *J Consult Clin Psychol*. 1997;65(1):79-85. doi:10.1037/0022-006X.65.1.79.
3. Jensen MD, Ryan DH, Apovian CM, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society [published correction appears in *Circulation*. 2014 Jun 24;129(25 Suppl 2):S139-40]. *Circulation*. 2014;129(25 Suppl 2):S102-S138. doi:10.1161/01.cir.0000437739.71477.ee.
4. Ryan DH, Yockey SR. Weight loss and improvement in comorbidity: differences at 5%, 10%, 15%, and over. *Curr Obes Rep*. 2017;6(2):187-194. doi:10.1007/s13679-017-0262-y.
5. Lopez-Minguez J, Gómez-Abellán P, Garaulet M. Timing of breakfast, lunch, and dinner. Effects on obesity and metabolic risk. *Nutrients*. 2019;11(11):2624. doi:10.3390/nu11112624.
6. Malik VS, Hu FB. The role of sugar-sweetened beverages in the global epidemics of obesity and chronic diseases. *Nat Rev Endocrinol*. 2022;18(4):205-218. doi:10.1038/s41574-021-00627-6.
7. Swift DL, McGehee JE, Earnest CP, et al. The effects of exercise and physical activity on weight loss and maintenance. *Prog Cardiovasc Dis*. 2018;61(2):206-213. doi:10.1016/j.pcad.2018.07.014.
8. Bellizzi A, van Baak MA, Battista F, et al. Effect of exercise training on weight loss, body composition changes, and weight maintenance in adults with overweight or obesity: An overview of 12 systematic reviews and 149 studies. *Obes Rev*. 2021;22 Suppl 4(Suppl 4):e13256. doi:10.1111/obr.13256.

The Ohio Cardiovascular and Diabetes Health Collaborative is funded by the Ohio Department of Medicaid and administered by the Ohio Colleges of Medicine Government Resource Center. The views expressed in this document are solely those of the authors and do not represent the views of the state of Ohio or federal Medicaid programs.

For more information head to [Cardi-OH.org](https://www.cardi-oh.org).

© 2023 Cardi-OH